


**Theoretical Physics**  
**Prof. Ruiz, UNC Asheville**  
**Chapter Q Homework. Laplace Transforms**

**HW-Q1. Laplace Transform.** Find the Laplace transform  $F(s)$  for the square pulse  $f(t)$  shown below by explicitly doing the Laplace transform integral. Then use the "derivative trick" for integration to obtain the Laplace transform  $G(s)$  of the ramp pulse  $g(t)$  from your result  $F(s)$  for the square pulse.



Finally, give  $F(1)$  and  $G(1)$  in terms of  $e$ , where  $e$  is the natural base.

**HW-Q2. Laplace Transform Shift Property.** Calculate the Laplace transform  $G(s)$  for

$$g(t) = t^n e^{-bt} \quad \text{two ways as described below, where } b > 0.$$

- a) Do the integral for the Laplace transform using the derivative trick.
- b) Use the shifting property: if  $g(t) = f(t)e^{at}$ , then  $G(s) = F(s - a)$ ,  $s > a$ .

**HW-Q3. Solving a Differential Equation.**




Image Courtesy David M. Harrison  
 Department of Physics  
 University of Toronto

Use Laplace transforms to solve the differential equation

$$m \frac{d^2 x}{dt^2} + b \frac{dx}{dt} + kx = 0$$

where you smack the block initially so that

$$x(0) = 0 \quad \text{and} \quad v(0) = A \sqrt{\frac{k}{m} - \frac{b^2}{4m^2}}.$$

Simply your math using these definitions:

$$\omega_0 = \sqrt{\frac{k}{m}}, \quad \beta = \frac{b}{2m}, \quad \omega^2 = \omega_0^2 - \beta^2.$$