

Open Book/Notes/Internet (90 minutes including scan and email return).

All work and related steps must be explicitly shown for full credit.

[10] 1. **Groups.** A group $G = \{a, b, c, d, e\}$ with identity e . For any $a \in G$ and $b \in G$, the following is true: $aba^{-1}b^{-1} = e$. Prove that the group is abelian.

[25] 2. **Integral.** Use a derivative trick to evaluate $\int_0^{\infty} xe^{-ax} \sin(kx) dx$, where $a > 0$,

starting from the integral result $\int_0^{\infty} e^{-ax} \cos(kx) dx = \frac{a}{a^2 + k^2}$. For 20 points max,

you may instead evaluate $\int_0^{\infty} xe^{-ax} \cos(kx) dx$. Commit to one for official credit.

[25] 3. **Waves.** Show that $\psi(x, t) = Ae^{-i(ax+bt)}$ satisfies the wave equation with an associated auxiliary equation that relates a , b , and the velocity v . What is this equation, in simplest form, that relates a , b , and v ?


[10] 4. **E&M.** If the second Maxwell equation was modified to be $\nabla \cdot \vec{B} = f(x, y, z) \neq 0$, explain in a sentence or two the new physics of this novel situation.

[30] 5. **Gas and Work.** A gas expands from point a to point b (see figure), where the

pressure varies as $P = \frac{k}{V^2}$ during the

expansion (k is a constant). What is the work done by the gas if the gas expands from $V_1 = V_0$ to $V_2 = 2V_0$ as shown in the figure, i.e., from point a to point b?

Give your answer in terms of P_0 and V_0 , where k does not appear in your answer.



1. **Groups.** Given: $aba^{-1}b^{-1} = e$. Multiply both sides by b on the right. Then $aba^{-1}b^{-1}b = eb \Rightarrow aba^{-1}e = b \Rightarrow aba^{-1} = b$. Multiply both sides by a on the right to get $aba^{-1}a = ba$, which leads to $abe = ba$ and finally $ab = ba$ (abelian).

2. **Integral.** $\int_0^{\infty} xe^{-ax} \sin(kx) dx$, where $a > 0$ using a derivative.

$$\int_0^{\infty} xe^{-ax} \sin(kx) dx = -\frac{d}{dk} \int_0^{\infty} e^{-ax} \cos(kx) dx = -\frac{d}{dk} \left(\frac{a}{a^2 + k^2} \right)$$

$$\int_0^{\infty} xe^{-ax} \sin(kx) dx = -\left[-\frac{a}{(a^2 + k^2)^2} \right] 2k = \frac{2ak}{(a^2 + k^2)^2}$$

3. **Waves.** The function $\psi(x, t) = Ae^{-i(ax+bt)} \Rightarrow \frac{\partial \psi}{\partial x} = -iaAe^{-i(ax+bt)} = -ia\psi$

$$\frac{\partial^2 \psi}{\partial x^2} = (-ia)^2 \psi = -a^2 \psi \quad \frac{\partial^2 \psi}{\partial t^2} = (-ib)^2 \psi = -b^2 \psi$$

Wave equation $\frac{\partial^2 \psi}{\partial x^2} = \frac{1}{v^2} \frac{\partial^2 \psi}{\partial t^2}$ is satisfied if $a^2 = \frac{b^2}{v^2} \Rightarrow v = \pm \frac{b}{a}$.

4. **E&M.** Given $\nabla \cdot \vec{B} = f(x, y, z) \neq 0$ would imply that magnetic charge exists, i.e.,

magnetic monopoles. Note that $\nabla \cdot \vec{E} = \frac{\rho}{\epsilon_0} \neq 0$ implies the existence of electric charge.

5. **Gas and Work.** Work $W = \int_{V_1}^{V_2} PdV = \int_{V_1}^{V_2} \frac{k}{V^2} dV = -k \frac{1}{V} \Big|_{V_1}^{V_2} = -k \frac{1}{V} \Big|_{V_0}^{2V_0}$

$$W = -k \left(\frac{1}{2V_0} - \frac{1}{V_0} \right) = \frac{k}{V_0} \left(1 - \frac{1}{2} \right) = \frac{k}{2V_0} \quad \text{Note that } 4P_0 = \frac{k}{V_0^2}.$$

Use $k = 4P_0V_0^2$ to obtain $W = \frac{k}{2V_0} = \frac{4P_0V_0^2}{2V_0} = 2P_0V_0$.

APPENDIX (Alternative Choice for Problem 2).

$$\int_0^{\infty} x e^{-ax} \cos(kx) dx = -\frac{d}{da} \int_0^{\infty} e^{-ax} \cos(kx) dx$$

$$\int_0^{\infty} x e^{-ax} \cos(kx) dx = -\frac{d}{da} \left(\frac{a}{a^2 + k^2} \right). \text{ Use the product rule.}$$

$$\int_0^{\infty} x e^{-ax} \cos(kx) dx = -\frac{1}{a^2 + k^2} - (-1) \frac{a}{(a^2 + k^2)^2} 2a$$

$$\int_0^{\infty} x e^{-ax} \cos(kx) dx = -\frac{1}{a^2 + k^2} + \frac{2a^2}{(a^2 + k^2)^2}$$

$$\int_0^{\infty} x e^{-ax} \cos(kx) dx = \frac{-(a^2 + k^2) + 2a^2}{(a^2 + k^2)^2} = \frac{a^2 - k^2}{(a^2 + k^2)^2}$$