

Theoretical Physics
Prof. Ruiz, UNC Asheville
Chapter N Homework. The Dirac Delta Function

HW-N1 and HW-N2. Probability Distribution and Moments. The n^{th} central moment for the probability distribution $P(x)$ is defined as

$$E[(x - \mu)^n] \equiv \int_{-\infty}^{\infty} (x - \mu)^n P(x) dx .$$

The "E" stands for expected value. Physicists like to use the term "expectation value" and use brackets. You will calculate some moments for the Gaussian centered on $x = 0$. **Find the zeroth, first, second (HW-N1), third, and fourth moments (HW-N2)** for

$$P(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{x^2}{2\sigma^2}} .$$

You **MUST NOT LOOK UP** any integrals, except the following one, and **YOU MUST NOT USE INTEGRATION BY PARTS**. Instead, from this integral already proven in class,

$$I = \int_{-\infty}^{\infty} e^{-\alpha x^2} dx = \sqrt{\frac{\pi}{\alpha}} ,$$

use the derivative trick to evaluate the following two integrals, which you will need.

$$I_1 = \int_{-\infty}^{\infty} x^2 e^{-\alpha x^2} dx \quad \text{and} \quad I_2 = \int_{-\infty}^{\infty} x^4 e^{-\alpha x^2} dx$$

After you evaluate the above integrals, choose the appropriate α for your problem.

HW-N3. Dirac Delta Function. Evaluate the following two integrals, showing all steps.

$$I_{k>0} = \int_{-\infty}^{\infty} f(x) \delta(kx) dx , \text{ where } k > 0$$

$$I_{k<0} = \int_{-\infty}^{\infty} f(x) \delta(kx) dx , \text{ where } k < 0$$

Hint: Let $z = kx$ and use what you know about the delta function from class.